

Please reference the tables below when considering how long to stop your anti-coagulant and injectable diabetic/weight loss medications before your procedure. Please also consult with your primary care doctor, your heart/lung specialist, or the prescribing physician for your anti-coagulant and diabetic medications for further clarification on the safety of stopping your medication (unless we've already done this for you). If your doctor denies you permission to stop your medication, please let your gastroenterologist know.

ANTI - COAGULANT MEDICATIONS	
Brilinta (ticagrelor)	Hold for 5 days prior
Coumadin (warfarin)	Hold for 5 days prior
Effient (prasugrel)	Hold for 7 days prior
Eliquis (apixaban)	Hold for 2-4 days (normal kidney function - 2 days)
Plavix (clopidogrel)	Hold for 5 days prior
Pradaxa (dabigatran)	Hold for 2-4 days (normal kidney function - 2 days)
Savaysa (edoxaban)	Hold for 1 day prior
Xarelto (rivaroxaban)	Hold for 2 day prior
Pletal (cilostazol)	Hold for 2 days prior
Aggrenox (aspirin & dipyridamole)	Hold for 7 days prior
Aspirin	Do not stop 81 or 325 mg. For 500 mg or more, STOP for 7 days

If you take an injectable GLP-1 medication below, please hold for 7 days prior to procedure. If you take an oral GLP-1 medication please hold the day before and the day of your procedure.

DIABETIC AND WEIGHT LOSS - INJECTABLE MEDICATIONS	
Byetta (exenatide)	Hold on day of procedure (twice daily injection)
Victoza (liraglutide)	Hold on day of procedure (once daily injection)
Tanzeum (albiglutide)	Hold for 7 days prior to procedure (once weekly injection)
Trulicity (dulaglutide)	Hold for 7 days prior to procedure (once weekly injection)
Lixumia (lixisenatide)	Hold on day of procedure (once daily injection)
Beinaglutide	Hold on day of procedure (three times daily injection)
Ozempic (semaglutide)	Hold for 7 days prior to procedure (once weekly injections)
Fu Laimel (peg-loxenate)	Hold for 7 days prior to procedure (once weekly injection)
Mounjaro (tirzepatide)	Hold for 7 days prior to procedure (once weekly injection)
Wegovy (semaglutide)	Hold for 7 days prior to procedure (once weekly injection)
Bydureon (exenatide)	Hold on day of procedure (twice daily injection)
Rybelsus (semaglutide)	Hold for 7 days prior to procedure (once weekly injection)
Soliqua (lixisenatide)	Hold on day of procedure (once daily injection)
Xultophy (Insulin degludec/liraglutide)	Hold on day of procedure (once daily injection)